

Plane Algebraic Curves – Problem Set 2

due Tuesday, May 12 at 10:15

- (1) (a) Find all singular points of the curve $F = (x^2 + y^2 - 1)^3 + 10x^2y^2 \in \mathbb{R}[x, y]$, and determine the multiplicities and tangents to F at these points.
- (b) Show that an irreducible curve F over a field of characteristic 0 has only finitely many singular points.
Can you find weaker assumptions on F that also imply that F has only finitely many singular points?

(2) Show:

- (a) If $F, G \in K[x_0, \dots, x_n]$ are polynomials such that $F \mid G$ and G is homogeneous, then F is homogeneous.
- (b) Every homogeneous polynomial in two variables over an algebraically closed field is a product of linear polynomials.

(3) By a *projective coordinate transformation* we mean a map $f: \mathbb{P}^n \rightarrow \mathbb{P}^n$ of the form

$$(x_0 : \dots : x_n) \mapsto (f_0(x_0, \dots, x_n) : \dots : f_n(x_0, \dots, x_n))$$

for linearly independent homogeneous linear polynomials $f_0, \dots, f_n \in K[x_0, \dots, x_n]$.

- (a) Let $P_1, \dots, P_{n+2} \in \mathbb{P}^n$ be points such that any $n+1$ of them are linearly independent in K^{n+1} , and in the same way let $Q_1, \dots, Q_{n+2} \in \mathbb{P}^n$ be points such that any $n+1$ of them are linearly independent. Show that there is a projective coordinate transformation f with $f(P_i) = Q_i$ for all $i = 1, \dots, n+2$.
- (b) Let F and G be two real smooth projective conics with non-empty set of points. Show that there is a projective coordinate transformation of \mathbb{P}^2 that takes F to G .
- (4) Let P be a point on an affine curve F . We say that P is a *cuspidal point* if $m_P(F) = 2$, there is exactly one tangent L to F at P , and $\mu_P(F, L) = 3$.
- (a) Give an example of a real curve with a cusp, and draw a picture of it.
- (b) If F has a cusp at P , prove that F has only one irreducible component passing through P .
- (c) If F and G have a cusp at P , what is the minimum possible value for the intersection multiplicity $\mu_P(F, G)$?